

On the $\Delta I = 1/2$ Rule in Holographic QCD

Thomas Hambye^{a,b}, Babiker Hassanain^a, John March-Russell^a, and Martin Schvellinger^a

^a*Rudolf Peierls Centre for Theoretical Physics, Department of Physics,
University of Oxford, 1 Keble Road, Oxford, OX1 3NP, UK*

^b*IFT-UAM/CSIC, Fac. Ciencias, Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain*

We study the $\Delta I = 1/2$ rule for kaon decays and the B_K parameter for $K^0 - \bar{K}^0$ mixing in a dual 5-dimensional holographic QCD model. We perform, in the chiral limit, computations of the relevant four-point current-current correlators, which depend upon self-interactions among the 5D bulk fields. Spontaneous chiral symmetry breaking (χ SB) is realized through boundary conditions on the bulk fields. Numerical results are analyzed in comparison with QCD, chiral perturbation theory (χ PT) and data, finding reasonable agreement with the experimental values of the g_8 and g_{27} parameters describing the $\Delta I = 1/2, 3/2$ decay channels.

I. INTRODUCTION

The gauge/gravity duality [1, 2] provides new tools to investigate non-perturbative QCD. Holographic dual models which allow the study of confinement and χ SB have been developed in essentially two different approaches: models based on string theory/supergravity in 10D [3], or phenomenologically inspired holographic dual models in 5D [4, 5, 6, 7, 8]. In the past year, a study of meson masses and decay constants in the 5D models have shown quite good agreement with data, at least for the leading meson excitations. These encouraging results are based on two-point current-current correlators, and in the 5D description the leading contributions to such correlators are obtained from bare 5D propagators. One can also compute n -point correlators at tree level; however, the fact that 5D interactions must be taken into account makes an important difference that we shall investigate here. The holographic calculation of four-point correlators is a non-trivial test for 5D dual models of non-perturbative QCD, with, perhaps, the most interesting aspect being the holographic computation of observables which are important in QCD, particularly the celebrated $\Delta I = 1/2$ rule for kaon decays. Since our calculation is the first within this framework, we focus upon the simplest AdS/QCD model.

II. THE $\Delta I = 1/2$ RULE, AND THE B_K MIXING PARAMETER

We start with a review of the relevant facts. Neglecting CP-violation there are two independent K^0 decays: $K^0 \rightarrow \pi^+\pi^-$ and $K^0 \rightarrow \pi^0\pi^0$. These two decays are combinations of $\Delta I = 1/2$ and $\Delta I = 3/2$ isospin amplitudes (A_0 and A_2 , respectively). Experimentally $ReA_0/ReA_2 = 22.2$, and the unexpected largeness of this ratio is the $\Delta I = 1/2$ rule. In the chiral limit, and at $\mathcal{O}(p^2)$ in the chiral expansion, these two amplitudes are expressed in terms of the g_8 and g_{27} parameters (see e.g. [9, 10]). In the following we will also discuss the related but simpler observable, $\hat{B}_K$, parameterizing $K^0 - \bar{K}^0$ mixing. Performing an OPE, its calcu-

lation reduces to that of the hadronic matrix element $\langle \bar{K}^0 | Q_{\Delta S=2} | K^0 \rangle (\mu) \equiv \frac{4}{3} f_K^2 M_K^2 B_K(\mu)$ of the four-quark left handed current operator $Q_{\Delta S=2} \equiv 4\eta_{\mu\nu} L_{sd}^\mu L_{sd}^\nu$ with $L_{q_1 q_2}^\mu \equiv \bar{q}_1 \gamma^\mu \frac{(1-\gamma_5)}{2} q_2$. Similarly, $g_{8,27}$ involve the $K \rightarrow \pi\pi$ matrix elements of two $\Delta S = 1$ operators four-quark left handed current operators (above the charm threshold), $Q_1 \equiv 4\eta_{\mu\nu} L_{sd}^\mu L_{uu}^\nu$ and $Q_2 \equiv 4\eta_{\mu\nu} L_{su}^\mu L_{ud}^\nu$.

The factorized contribution for these matrix elements is well known; the unfactorized contribution is more controversial. In the chiral limit the leading N_c unfactorized contributions reduce to the calculation of integrals in Q^2 (with Q the Euclidean momentum flowing between the left handed currents) of two distinct four-point current correlators [10, 11]. For $\hat{B}_K \equiv B_K(\mu) \cdot C_{\Delta S=2}(\mu)$,

$$B_K(\mu) = \frac{3}{4} - \frac{3}{4} \frac{1}{32\pi^2 F_\pi^2} \int dQ^2 W_{LRLR}(Q^2)(\mu), \quad (1)$$

$$W_{LRLR}(Q^2) = -\frac{Q^2}{3F_\pi^2} \eta_{\alpha\beta} \eta_{\mu\nu} \int \frac{d\Omega_q}{4\pi} W_{LRLR}^{\mu\alpha\nu\beta}(q),$$

$$W_{LRLR}^{\mu\alpha\nu\beta}(q) = \lim_{l \rightarrow 0} i^3 \int d^4x d^4y d^4z e^{iqx + il(y-z)} \langle 0 | T \{ L_{sd}^\mu(x) R_{ds}^\alpha(y) L_{sd}^\nu(0) R_{ds}^\beta(z) \} | 0 \rangle |_{\text{conn}}, \quad (2)$$

while g_8 and g_{27} are given by

$$g_8(\mu) = z_1(\mu) \left(\frac{4B_K(\mu)}{5} - 1 \right) + \quad (3)$$

$$z_2(\mu) \left(1 - \frac{8B_K(\mu)}{15} - \int dQ^2 \frac{W_{LLRR}(Q^2)}{4\pi^2 F_\pi^2}(\mu) \right),$$

$$g_{27}(\mu) = \left(z_1(\mu) + z_2(\mu) \right) \frac{4B_K(\mu)}{5}. \quad (4)$$

In the above, $C_{\Delta S=2}(\mu)$ and $z_{1,2}(\mu)$ are the Wilson coefficients of $Q_{\Delta S=2}$ and $Q_{1,2}$, $R_{q_1 q_2}^\mu \equiv \bar{q}_1 \gamma^\mu \frac{(1+\gamma_5)}{2} q_2$, and

$$W_{LLRR}(Q^2) = -\frac{Q^2}{3F_\pi^2} \eta_{\alpha\beta} \eta_{\mu\nu} \int \frac{d\Omega_q}{4\pi} W_{LLRR}^{\mu\nu\alpha\beta}(q),$$

$$W_{LLRR}^{\mu\nu\alpha\beta}(q) = \lim_{k \rightarrow 0} i^3 \int d^4x d^4y d^4z e^{iqx + ik(y-z)} \langle 0 | T \{ L_{su}^\mu(x) L_{ud}^\nu(0) R_{du}^\alpha(y) R_{us}^\beta(z) \} | 0 \rangle |_{\text{conn}}. \quad (5)$$

III. THE HOLOGRAPHIC QCD MODEL

To compute the four-point correlators through the gauge/gravity correspondence, we consider the $L_{\bar{q}_1 q_2}^\mu$ and $R_{\bar{q}_1 q_2}^\mu$ 4D quark currents. By virtue of the AdS/CFT correspondence, these operators couple to the boundary values of 5D gauge fields $L^\mu(x^\nu, z)$ and $R^\mu(x^\nu, z)$, and the 5D gauge fields are massless. The 5D action is a $SU(3)_L \times SU(3)_R$ Yang-Mills theory

$$-\frac{M_5}{4} \int dz d^4x \sqrt{g} (L_{MN}^a L^{a, MN} + R_{MN}^a R^{a, MN}), \quad (6)$$

where M_5 is an undetermined mass scale, and $M = (\mu, 5)$. The fifth coordinate z runs from L_0 to L_1 , which are the positions of the UV and IR branes respectively. It is possible to set $L_0 = 0$ as in Refs.[4, 5] as our results are smooth as $L_0 \rightarrow 0$. We use the AdS_5 metric $ds^2 = a(z)^2(\eta_{\mu\nu} dx^\mu dx^\nu - dz^2)$, where $a(z)^2 = (L/z)^2$ and $\eta_{\mu\nu}$ is given by $(+, -, -, -)$. The field strengths are $L_{MN} = \partial_M L_N - \partial_N L_M - i[L_M, L_N]$, where $L_M = L_M^a T^a$, and similarly for R_{MN} . We normalise $\text{tr}(T^a T^b) = \delta^{ab}$, and define the vector and axial gauge bosons as $V_M = (L_M + R_M)/\sqrt{2}$ and $A_M = (L_M - R_M)/\sqrt{2}$, respectively.

The holographic dual of this 5D theory is a 4D theory with three quark flavours and a global $SU(3)_L \times SU(3)_R$ chiral symmetry [4, 5, 6]. We work in the chiral limit and include the effect of the spontaneous χ SB via the IR boundary condition on the axial vector fields [5, 6]. This is clearly a rather severe limit, yet the calculational simplification it affords makes it worthwhile. (The effects of spontaneous and explicit χ SB could be treated more precisely by the inclusion of a bulk scalar in the bi-fundamental representation of the gauge group [4, 5].)

A. Propagators and 5D Interactions

By the holographic correspondence, the UV boundary values of V_μ and A_μ act as classical sources coupled to the 4D vector and axial global symmetry currents. Moreover, the 5D action evaluated on the solutions of the equations of motion (EOM) of the bulk fields defines the generating functional for current-current correlators in the 4D theory. We therefore need to solve the EOM with given UV boundary values of the fields, and substitute back into the action. The EOM are non-linear, so we must solve them iteratively. This is the origin of Witten diagram construction for calculating the n -point Greens functions of operators in the dual theory [2]. The iterative solution of the EOM requires the bulk-to-bulk and bulk-to-boundary propagators for the gauge fields in the 5D theory.

We work in the unitary gauge in both the $SU(3)_V$ and $SU(3)_A$ sectors. V_5 is eliminated completely in this gauge if we impose null Dirichlet boundary conditions for it on both branes. We impose warped Neumann boundary conditions on the A_5 field (i.e. $\partial_z(aA_5) = 0$ on both

branes). In this gauge A_5 is a physical field because it has a zero mode which cannot be gauged away. The V_μ and A_μ propagators have both transverse and longitudinal parts in this gauge. Following Ref.[12] we have

$$\langle V^\mu V^\nu \rangle = -iG_p^V(z, z') P_T^{\mu\nu} - iG_0^V(z, z') P_L^{\mu\nu}, \quad (7)$$

where $P_T^{\mu\nu} = \eta^{\mu\nu} - \frac{p^\mu p^\nu}{p^2}$ and $P_L^{\mu\nu} = \frac{p^\mu p^\nu}{p^2}$. The A_μ propagator has a similar form, and both solve

$$\left(\partial_z^2 - \frac{1}{z}\partial_z + p^2\right) G_p^{V,A}(z, z') = \frac{z\delta(z-z')}{M_5 L}. \quad (8)$$

The boundary conditions on $G_p^V(z, z')$ are Dirichlet on the UV brane, $G_p^V(z, z')|_{z=L_0} = 0$, and Neumann on the IR brane, $\partial_z G_p^V(z, z')|_{z=L_1} = 0$. We impose Dirichlet boundary conditions on both branes in the axial sector $G_p^A(z, z')|_{z=L_0, L_1} = 0$, to encode χ SB. The bulk-to-boundary propagator is defined as:

$$\langle V^\mu V^\nu \rangle \Big|_{\partial ADS} (z') = -\frac{M_5 L}{z} \partial_z \langle V^\mu V^\nu \rangle \Big|_{z=L_0}, \quad (9)$$

(with a similar equation for A_μ), where

$$\langle V^\mu V^\nu \rangle \Big|_{\partial ADS} (z') = -iK_p^V(z') P_T^{\mu\nu} - iK_0^V(z') P_L^{\mu\nu}, \quad (10)$$

$$K_p^{V,A}(z') = -\frac{M_5 L}{z} \partial_z G_p^{V,A}(z, z')|_{z < z'} \Big|_{z=L_0}, \quad (11)$$

and similarly for $K_0^{V,A}(z')$. The solutions are given by:

$$G_p^{V,A}(z, z')_{z < z'} = \pi z z' \times \frac{[A\mathcal{J}_1(pz) + B\mathcal{Y}_1(pz)][C\mathcal{J}_1(pz') + D\mathcal{Y}_1(pz')]}{2M_5 L(AD - BC)}, \quad (12)$$

$z > z'$ implies the exchange $z \leftrightarrow z'$. For the vector sector the coefficients are $A = -\mathcal{Y}_1(pL_0)$, $C = -\mathcal{Y}_0(pL_1)$, $B = \mathcal{J}_1(pL_0)$ and $D = \mathcal{J}_0(pL_1)$; for A_μ we obtain $A = -\mathcal{Y}_1(pL_0)$, $C = -\mathcal{Y}_1(pL_1)$, $B = \mathcal{J}_1(pL_0)$ and $D = \mathcal{J}_1(pL_1)$. In the zero momentum limit:

$$G_0^V(z, z')_{z < z'} = -\frac{1}{2M_5 L}(z^2 - L_0^2), \quad (13)$$

($z > z'$ again swaps z with z'). For the axial sector

$$G_0^A(z, z')_{z < z'} = -\frac{1}{2M_5 L} \frac{(z^2 - L_0^2)(z'^2 - L_1^2)}{L_0^2 - L_1^2}, \quad (14)$$

and swapping z and z' for $z > z'$. The bulk-to-boundary propagators are then

$$K_p^{V,A}(z') = -\frac{z'}{L_0} \frac{C\mathcal{J}_1(pz') + D\mathcal{Y}_1(pz')}{AD - BC}, \quad (15)$$

while $K_0^V(z') = 1$, and $K_0^A(z') = \frac{z'^2 - L_1^2}{L_0^2 - L_1^2}$. The physical condition of 4D current conservation eliminates the




FIG. 1: The three basic structures of the 36 individual Witten diagrams: X-diagrams involving the 4-boson vertex, Y-diagrams involving two 3-boson vertices, and Y diagrams where A_5 propagates. The exterior circle represents the (UV) boundary of the AdS_5 space, the solid lines are boundary-to-bulk or bulk-to-bulk (axial or vector) gauge propagators, and the dotted line is a bulk-to-bulk A_5 propagator.

longitudinal piece of the bulk-to-boundary propagators. The positions of the poles of K_p^V and K_p^A are the masses of the vector and axial-vector mesons, respectively. This is the usual association of the masses of the Kaluza-Klein modes in 5D with the masses of the bound states of the dual theory [13]. Note that this tree-level treatment of the 5D model incorrectly predicts that the mass of the n 'th excited resonance scales like n (for $n \gg 1$) as opposed to the $\sqrt{n}$ Regge behaviour. This failing is to be expected, as the simple 5D theory here presented is invalid in the far UV [14].

We also note that the current-current correlator in the axial sector as $p \rightarrow 0$ is $\Pi_A(p^2)|_{p=0} = F_\pi^2 = 2M_5L/(L_1^2 - L_0^2)$ [4, 5, 6]. The A_5 propagator in this gauge is $\langle A_5 A_5 \rangle = -iG_p^5(z, z')$, where $G_p^5(z, z')$ is the limit as $\xi \rightarrow \infty$ of the solution of

$$\left(\xi \partial_z^2 - \xi \frac{1}{z} \partial_z + \xi \frac{1}{z^2} + p^2 \right) G_p^5(z, z') = -\frac{z \delta(z - z')}{M_5 L}, \quad (16)$$

with the boundary conditions $\partial_z(aG_p^5(z, z'))|_{z=L_0, L_1} = 0$, giving

$$G_p^5(z, z') = \frac{-zz'}{M_5 L(L_1^2 - L_0^2)} \frac{1}{p^2}. \quad (17)$$

The interactions are the three-gauge boson and four-gauge boson vertices derived from the 5D action Eq.(6).

B. Four-point current-current correlators

In order to calculate the four-point current-current correlators we have to solve tree level four-point Witten diagrams which contain either one four-gauge field vertex or two three-gauge field vertices. Note that certain 5d Witten diagrams do not contribute to the four-point functions (2) and (5) because they do not preserve the flavor structure of the 4d theory.

We find that the group theory structure is a common factor in all diagrams, so denoting by Σ the sum of all diagrams with group factors amputated we have $\Sigma =$

$\Sigma_X + \Sigma_{A_5} + \Sigma_Y$, with

$$\Sigma_X = 3iM_5L \int \frac{dz}{z} ([K_0^{V^2} + K_0^{A^2}][K_p^{V^2} + K_p^{A^2}] - 4K_0^V K_0^A K_p^V K_p^A) \quad (18)$$

$$\Sigma_{A_5} = -\frac{3i}{2}(M_5L)^2 \int \frac{dz}{z} \frac{dz'}{z'} G_p^5(z, z') A'(z, z') \quad (19)$$

where $A'(z, z')$ is a function whose explicit form is known. As for the Y-diagrams, the integrations are more involved, but the sum can be written as $\Sigma_Y = Y_1 + Y_2 + Y_3 + Y_4 - 2Y_5 - 2Y_6$, where

$$Y_i = -3i \left(\frac{M_5L}{\sqrt{2}} \right)^2 p^2 \int \frac{dz}{z} \frac{dz'}{z'} K_p^{a_i}(z) G_0^{b_i}(z, z') K_p^{c_i}(z') K_0^{d_i}(z) K_0^{e_i}(z') \quad (20)$$

with $\{abcde\} = \{VVVVV\}, \{VAVAA\}, \{AAAVV\}, \{AVAAA\}, \{AAVVV\}, \{AVVAV\}$ for $i = 1, \dots, 6$. Performing the Wick rotation to Euclidean space, we obtain:

$$W_{LRLR}(Q^2) = \frac{iQ^2}{6F_\pi^2} \Sigma(p = iQ) = -2W_{LLRR}(Q^2). \quad (21)$$

Thus we find that the ratio of the two correlators is independent of the integration in z -space, and is solely due to their different $SU(3)_L \times SU(3)_R$ symmetry factors. The integrals can be done analytically with finite values for L_1 and L_0 , but our observables have a smooth $L_0 \rightarrow 0$ limit, and much simpler expressions arise if we set $L_0 = 0$ giving:

$$\begin{aligned} \Sigma(Q) = 3iM_5L & \left[\frac{32}{Q^6 L_1^6} - \frac{16}{5Q^4 L_1^4} - \frac{69}{60I_1^2} + \frac{2}{5Q^2 L_1^2 I_1^2} \right. \\ & + \frac{69}{60I_0^2} - \frac{3}{15Q^2 L_1^2 I_0^2} + \frac{8}{5Q^4 L_1^4 I_0^2} \\ & \left. + \frac{32}{Q^6 L_1^6 I_0^2} - \frac{64}{Q^6 L_1^6 I_0} + \frac{1}{QL_1 I_0 I_1} \right], \quad (22) \end{aligned}$$

where $I_{0,1} \equiv I_{0,1}(QL_1)$ are modified Bessel functions of zeroth and first order respectively.

IV. RESULTS AND DISCUSSION

The three input parameters of our model are M_5L , L_0 and L_1 . From the expressions in Section II.A one can easily compute the boundary-to-boundary propagator in the vector sector, and match its large Euclidean momentum behaviour to that known from perturbative QCD deriving the relation[4, 5]

$$M_5L = \frac{N_c}{12\pi^2}. \quad (23)$$

One is then left with the two dimensionful parameters, $1/L_0$ giving the UV cutoff scale, and $1/L_1$ setting the

IR scale. As mentioned before, one problem of the simple 5D holographic model presented is that it does not correctly reproduce the spectrum of meson resonances in the regime above approximately 1500 MeV. In addition, in computing the physically relevant observables g_8 and g_{27} one must in practice choose a scale μ at which to evaluate the Wilson coefficients $z_1(\mu)$ and $z_2(\mu)$. One simple choice we make in the following is to take $1/L_0 = 1500$ MeV, imposing this as an upper hard cut-off when we evaluate the Q^2 integrals in Eqs.(1)-(5), and identifying this hard cut-off with the short distance renormalisation scale μ . Although this matching procedure is quite crude, Refs. [15] have shown that such a procedure captures the dominant contributions. In a later longer publication we investigate in some detail other more sophisticated matching procedures. With respect to the χ PT and perturbative QCD calculations of [15] and [10, 11] of the B_K and $\Delta I = 1/2$ rule observables, our model, if valid, presents the interesting possibility of calculating the intermediate Q^2 contribution around 0.1-2GeV² directly, rather than fitting it from an interpolation between the χ PT low Q^2 regime [15] and of the OPE high Q^2 regime [10, 11].

One immediate check on our calculation follows from the fact that the most general form for the Greens function $W_{LRLR}(Q^2)$ in the large N_c limit can be written in terms of the masses, M_i , and residues of the meson resonances [10, 11]

$$\sum_{i=1}^{\infty} \left(\frac{\alpha_i}{(Q^2 + M_i^2)} + \frac{\beta_i}{(Q^2 + M_i^2)^2} + \frac{\gamma_i}{(Q^2 + M_i^2)^3} \right), \quad (24)$$

and similarly for W_{LLRR} with residues α'_i , β'_i and γ'_i . We have explicitly checked that our results for $W_{LRLR}(Q^2)$ and $W_{LLRR}(Q^2)$ agree with this general form.

We now turn to the comparison of our results with the experimental data. Given Eq.(23) and our choice of $L_0 = 1/1500$ MeV⁻¹, the one remaining free parameter of our model is L_1 . The data to which this must be fitted are, most importantly, the pion decay constant, F_π , the ρ vector-meson mass, m_ρ , the a_1 axial-vector meson mass m_{a_1} , and the g_8 and g_{27} (or B_K) parameters. In the low- Q^2 regime chiral perturbation theory calculations [10, 11] give the behaviour of the correlation functions

$$\begin{aligned} W_{LRLR}(Q^2) &= 6 - 24(2l_1 + 5l_2 + l_3 + l_9) \frac{Q^2}{F_\pi^2} + \dots \\ W_{LLRR}(Q^2) &= -\frac{3}{8} + (-\frac{15}{2}l_3 + \frac{3}{2}l_9) \frac{Q^2}{F_\pi^2} + \dots \end{aligned} \quad (25)$$

with l_i the standard chiral-Lagrangian coefficients. In our fit procedure we use the χ PT results, Eq.(25), in the integrals over Q^2 below $1/L_1$, and the results of the holographic calculation, Eqs.(21) and (22) in the regime $1/L_1$ to $1/L_0$.

In terms of $M_5 L$, L_1 and L_0 the predictions of the 5D

holographic model for F_π and m_ρ and m_{a_1} are [4, 5, 6]

$$F_\pi^2 \approx \frac{2M_5 L}{L_1^2 - L_0^2} \quad (26)$$

and in a good approximation in the range of interest

$$m_\rho \approx \frac{2.12}{L_1} \frac{(L_1 - 0.282L_0)}{(L_1 - L_0)} \quad (27)$$

$$m_{a_1} \approx \frac{3.38}{L_1} \frac{(L_1 - 0.085L_0)}{(L_1 - L_0)}. \quad (28)$$

(We use a precise interpolation to the exact result for our fits.) A non-trivial success of the 5D holographic QCD model is that in the $L_0 = 0$ limit (to which $1/L_0 = 1500$ MeV is a good approximation) the ratio $m_{a_1}/m_\rho \approx 1.6$ is very well reproduced independently of the values of $M_5 L$ and L_1 , as already noted in [4, 5].

Numerically performing the Q^2 integrations in Eqs.(1), (4), and (5), as a function of $1/L_1$ and utilizing, as appropriate, the leading order values of the Wilson coefficients z_1 and z_2 (taken from [16] with $\Lambda_{\overline{MS}}^{(4)} = 325$ MeV, and $\mu = 1500$ MeV) we can fit the final parameter L_1 to the full set of observables. The $\Delta I = 1/2, 3/2$ data requires $g_8|_{obs} = 5.1$ and $g_{27}|_{obs} = 0.29$, although the values to be explained are modified to $g_8|_{obs} = 3.3$ and $g_{27}|_{obs} = 0.23$ when one takes into account the enhancement already provided by the calculated $\mathcal{O}(p^4)$ chiral corrections [9]. Fitting to g_8 and g_{27} and the values of F_π , m_ρ and m_{a_1} leads to the result

$$L_1^{-1} \approx 302 \text{ MeV} \quad (29)$$

which implies (in all cases normalized to the data for ease of understanding)

$$\begin{aligned} \frac{m_\rho|_{th}}{m_\rho|_{obs}} &\approx 1.00, \quad \frac{m_{a_1}|_{th}}{m_{a_1}|_{obs}} \approx 1.04, \quad \frac{F_\pi|_{th}}{F_\pi|_{obs}} \approx 0.80, \\ \frac{g_8|_{th}}{g_8|_{obs}} &\approx 0.49, \quad \frac{g_{27}|_{th}}{g_{27}|_{obs}} \approx 1.39 \end{aligned} \quad (30)$$

and a value of $B_K(1500 \text{ MeV}) \approx 0.54$ ($\hat{B}_K \approx 0.76$ using the Wilson coefficient value $C_{\Delta S=2}(1500 \text{ MeV}) = 1.42$, see e.g. Ref. [11]). Considering the relative crudity of the model and the use of the large- N_c expansion of QCD this is a reasonable fit to the data and we find this result very encouraging. We emphasise that L_1 and L_0 are the only free parameters in this fit to the five observables of equation Eq.(30). It is noteworthy that in the expected domain of validity of the model (i.e. in the Q^2 intermediate region), the 5D interactions induce a large increase of g_8 and a suppression for g_{27} .

On the other hand the holographic model predicts the low- Q^2 behaviour of W_{LRLR} and W_{LLRR} to be identical in form to χ PT but with disagreeing numerical values

$$W_{LLRR} = -\frac{W_{LRLR}}{2} = -\frac{1}{4} + \frac{M_5 L}{4} \frac{29}{24} \frac{Q^2}{F_\pi^2} + \dots \quad (31)$$

It is possible that this failing is due to the simplistic truncation of the AdS space in the IR. Alternatively, the fault might be with our approximations, particularly the treatment in the 5D model of χ SB.

It would be interesting to include a bi-fundamental bulk scalar field of mass-squared $-3/L^2$ associated with the $\bar{q}q$ operator, as in the models [4, 5], and a massless singlet scalar associated with the glueball states generated by the gluon field operator $G_{\mu\nu}G^{\mu\nu}$, to see whether their inclusion leads to an even better fit to the full set of observables.

Acknowledgments

This work has been supported by PPARC Grant PP/D00036X/1, and by the ‘Quest for Unification’ network, MRTN 2004-503369. The work of BH was supported by the Clarendon Fund and the one of TH by a Ramon y Cajal contract of the Ministerio de Educación y Ciencia.

-
- [1] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
 - [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Phys. Lett. B* **428**, 105 (1998). E. Witten, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
 - [3] M. Kruczenski et al, *JHEP* **0307**, 049 (2003); *JHEP* **0405**, 041 (2004). T. Sakai and S. Sugimoto, *Prog. Theor. Phys.* **113**, 843 (2005). J. Babington et al, *Phys. Rev. D* **69**, 066007 (2004); *Fortsch. Phys.* **52**, 578 (2004). S. Hong, S. Yoon and M. J. Strassler, *hep-ph/0501197*. G. F. de Teramond and S. J. Brodsky, *Phys. Rev. Lett.* **94**, 201601 (2005).
 - [4] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, *Phys. Rev. Lett.* **95**, 261602 (2005).
 - [5] L. Da Rold and A. Pomarol, *Nucl.Phys. B* **721** (2005) 79; L. Da Rold and A. Pomarol, *JHEP* **0601**, 157 (2006).
 - [6] J. Hirn and V. Sanz, *JHEP* **0512**, 030 (2005); J. Hirn and V. Sanz, *arXiv:hep-ph/0510023*.
 - [7] K. Ghoroku and M. Yahiro, *Phys. Lett. B* **604**, 235 (2004); K. Ghoroku, T. Sakaguchi, N. Uekusa and M. Yahiro, *Phys. Rev. D* **71**, 106002 (2005); K. Ghoroku, N. Maru, M. Tachibana and M. Yahiro, *Phys. Lett. B* **633**, 602 (2006).
 - [8] E. Katz, A. Lewandowski and M. D. Schwartz, *hep-ph/0510388*.
 - [9] J. Kambor, J. Missimer and D. Wyler, *Phys. Lett. B* **261**, 496 (1991); E. Pallante, A. Pich and I. Scimemi, *Nucl.Phys. B* **617**, 441 (2001); J. Bijnens, P. Dhonte and F. Persson, *Nucl.Phys. B* **648**, 317 (2003).
 - [10] T. Hambye, S. Peris and E. de Rafael, *JHEP* **0305**, 027 (2003).
 - [11] S. Peris and E. de Rafael, *Phys. Lett. B* **508**, 117 (2001).
 - [12] L. Randall and M. D. Schwartz, *JHEP* **0111** (2001) 003.
 - [13] N. Arkani-Hamed, M. Porrati and L. Randall, *JHEP* **0108**, 017 (2001).
 - [14] Very recently, it has been shown that in a theory with linear confinement Regge behaviour imposes a strong constraint on the infrared dynamics of the holographic QCD model, see A. Karch, E. Katz, D. T. Son and M. A. Stephanov, *arXiv:hep-ph/0602229*.
 - [15] W. Bardeen, A. Buras and J.-M. Gérard, *Nucl.Phys. B* **293**, 787 (1987); T. Hambye et al, *Phys.Rev.* **D58**, 014017 (1998); *Eur.Phys.J. C* **10** (1999) 271; J. Bijnens and J. Prades, *Nucl.Phys. B* **444**, 523 (1995); *JHEP* **023**, 9901 (1999).
 - [16] G. Buchalla, A.J. Buras and M.E. Lautenbacher, *Rev.Mod.Phys.* **68**, 1125 (1996).